

EARTHWORMS IN THE HIMALAYA AND WESTERN GHATS REGION OF INDIA: A REVIEW

MADHAB C. DASH* AND K. G. SAXENA¹

Odisha Environment Appellate Authority,
Nilakantha Nagar, Bhubaneswar - 751 012

¹School of Environmental Sciences, Jawaharlal Nehru University, New Delhi – 100 067

E-mail: kgsaxena@jnu.ac.in

KEY WORDS

Soil biodiversity,
Mountainous region,
Soil macro-fauna,
Earthworm

Received on :

22.12.2011

Accepted on :

20.02.2012

***Corresponding
author**

ABSTRACT

Soil biodiversity is one of the most challenging areas. Our understanding about belowground biodiversity, especially the soil faunal component and its functions in mountainous ecosystems are not as strong as that of other ecosystems and on aboveground biodiversity. Earthworms help faster nutrient release from the organic substrates. This paper provides a review of the available knowledge on earthworm diversity, community structure and abundance in the Himalayas and the Western Ghats, the regions distinguished as global biodiversity hotspots based on the assessment of aboveground diversity.

INTRODUCTION

Mountain regions of the world are fascinating as they cover a wide range of ecological diversity over smaller areas because of elevation effect and settlement of a several local communities maintaining natural ecosystems together with the managed ones, with landscape management practices varying among local communities isolated by terrain and linguistic barriers. In recent times ecotourism to mountainous regions is not only an economic activity but demands conservation of traditional as well as new ecosystems that attract tourists. Sustainable land use and resource management are key requirements for an area to get recognition as a preferred spot of ecotourism. Increasing demand for organic food and persistence of traditional organic agriculture as patches in the matrix of natural ecosystems in marginal mountain areas make them a prospective area of ecotourism. Soil biodiversity is key to sustainable organic farming (Ramkarishnan *et al.*, 2005) and earthworms are the most dominant component of soil biota in terms of biomass and crucial for maintaining soil fertility (Dash, 1978, Senapati and Dash, 1981, Julka and Paliwal, 2005a and b; Dash *et al.*, 2009; Bhadauria *et al.*, 2012; Dash, 2012). This paper is an attempt to review the information available on diversity and functions of earthworms in the Himalaya and Western Ghats region of India, the areas distinguished globally for their highly valuable biodiversity and ecosystem services, and to identify knowledge gaps to be addressed in future research.

Distribution and diversity of earthworms: national survey

INVENTORY India is spread over an area of 3,287,797 km²

(2.4% area of the World), covering a wide range of physiographic, climatic and land use/cover types. Among the 15 agro-climatic regions identified in the country, (Fig. 1) four regions viz., Eastern Himalaya, North-east Ranges, Western Ghats and Western Himalayas figure among the global biodiversity hotspots. However, these hotspots identified based on extremely high endemic vascular plant species richness (presence of at least 0.5% or 1500 endemic vascular plant species) and vegetation degradation (loss of at least 70% primary vegetation) may not necessarily have high diversity of other plant and animal taxa (Myers, 2000; Kareiva and Marvier, 2003).

Geological histories, socio-cultural and economic conditions differ within as well as between hotspots. Thus, settled agriculture on terraced slopes constitutes the predominant agricultural land use in the western and central Himalaya and shifting/slash-burn agriculture on natural slopes in the eastern Himalaya and its extension ranges. The Himalayas is a creation of modern plate tectonic forces but not the Western Ghats, an area that has never been submerged under sea. In contrast to high-input commercial agriculture that dominates in the Indo-Gangetic plains, subsistence low-input traditional farming is widespread in the mountain regions (Bhadauria and Ramakrishnan, 2005; Senapati *et al.*, 2005; Chaudhuri *et al.*, 2008).

Zoological Survey of India has been involved with inventorying of soil fauna diversity for a long period of time, with survey efforts focused largely on presence/absence of different taxa in different environments. Nevertheless, the efforts of this organization devoted exclusively for survey and inventorying

of the faunal wealth of the country are augmented by many researchers in other research and development institutions. Nine families of earthworms with 69 genera and 418 species have been reported from India. On the basis of available data, the Western Ghats and West Coastal Plains would stand out as the regions with the highest level of earthworm species richness followed by Eastern Himalayan Region, Southern Plateau, Western Himalayan Region, Eastern Coastal Plains and Eastern Ghats, Gangetic Plains, Gujarat plains, Islands, Western dry Regions and transgangetic regions (Julka and Mukherjee, 1984; Julka and Paliwal, 2005a and b; Dash and Dash, 2008). The Western Ghats region is home to 53% species known from India compared to 26% and 12% in the case of the Eastern Himalaya and the Western Himalaya, respectively. *Drawida* (38 species) is the most species rich genus followed by *Megascolex* (30 species) in the Western Ghats, *Perionyx* (33 species) followed by *Drawida* (14 species) in the Eastern Himalaya, and *Perionyx* and *Amyntas* (4 species in each Genus) in the Western Himalaya. The Western Ghats harbour 193 native species compared to 85 in the Eastern Himalaya and 22 in the Western Himalaya, though the three regions do not differ much in terms of number exotic species (25-26). Among the native peregrine species associated with agroecosystems, *Octochaetona palniensis* is confined to the Western Ghats and *Lenogaster pusillus* and *Eutyphoeus* spp. to the Western Himalaya. Among the exotics associated with agroecosystems, *Drawida japonica* was able to establish only in the Western Himalaya, while *Dichogaster affinis* and *Pontoscolex corethrurus* could establish in Eastern Himalaya and Western Ghats region but not in the Western Himalaya (Table 1).

Earthworm communities and populations

Studies in village landscapes in central/western Himalaya (Bhadoria *et al.*, 2000, Sinha *et al.*, 2003) showed occurrence of three species in pasture soils compared to eight species in other land-uses, with Exotic *Amyntas corticis* being the most common species. Biological invasion was observed in both early and late successional stages.

Bhadoria and Ramakrishnan (2005) found three native species, viz., *Tonoscolex horai*, *Drawida assamensis* and *Perionyx* sp. in the primary forest in north-east India. Conversion of primary forest for slashes and burn agriculture resulted in the loss of two native species and colonization of the disturbed area by another native species viz., *Nelloscolex strigosus* and exotic *Amyntas corticis*. Although richness of native species increased and the native species were able to coexist with the exotic species during secondary succession after abandonment of cultivation, complete restoration of native earthworm species assemblage did not occur. A similar trend was observed in central/western Himalayan region, though there were differences in species found in the western and eastern Himalaya (Bhadoria *et al.*, 2012).

Chaudhuri *et al.* (2008) found that conversion of primary forests to rubber plantations in Tripura led to dominance of the exotic-endogeic *Pontoscolex corethrurus* as well as a change in the functional status of this species, from endogeic in the primary forests to endo-aneic and endo-epigeic in plantations. In general the change in land use pattern or loss

of primary forest followed establishment of some exotic species and also replacement of some natives by other native species.

Epigeic and anecic species, such as *Dichogaster bolau*, *Drawida willsi*, *Perionyx excavatus*, *Perionyx sansibaricus*, *Ramiella* sp. and *Lampito mauritii* are widely distributed and valued for their use in vermitechnology in Western and Eastern

Table 1: Native peregrine and exotic earthworm species associated with agroecosystems in the Himalaya and the Western Ghats region of India (based on Julka and Paliwal, 2005a & b and personal communication with Julka; A, absent; P, present)

Native peregrine species	Western Himalaya	Eastern Himalaya	Western Ghats
<i>Lampito mauritii</i>	A	P	P
<i>Perionyx excavatus</i>	P	P	P
<i>Perionyx sansibaricus</i>	P	A	P
<i>Octochaetona beatrix</i>	P	A	P
<i>Octochaetona surensis</i>			
<i>Octochaetona palniensis</i>	A	A	P
<i>Lenogaster pusillus</i>	P	A	A
<i>Ramiella bishambari</i>	P	A	P
<i>Eutyphoeus incommodus</i>	P	A	A
<i>Eutyphoeus michaelsoni</i>	P	A	A
<i>Eutyphoeus waltoni</i>	P	A	A
<i>Drawida willsi</i>	A	A	A
<i>Drawida calebi</i>	A	A	A
<i>Drawida nepalensis</i>	P	P	A
<i>Thaonia gracilis</i>	P	A	A
Exotic species			
<i>Dichogaster affinis</i>	A	P	P
<i>Dichogaster bolau</i>	P	P	P
<i>Amyntas alexandri</i>	P	P	P
<i>Amyntas corticis</i>	P	P	P
<i>Amyntas morrisi</i>	P	P	P
<i>Metaphire houlleti</i>	P	P	P
<i>Metaphire posthuma</i>	P	P	P
<i>Polyphreirima eleongata</i>	A	P	P
<i>Drawida japonica</i>	P	A	A
<i>Pontoscolex corethrurus</i>	A	P	P
<i>Ocnodrilus occidentalis</i>	P	A	P
<i>Allobophora parva</i>	P	P	P
<i>Aporrectodea cal. trapezoides</i>	P	P	P
<i>Aporrectodea rosea rosea</i>	P	P	P
<i>Eisenia fetida</i>	P	P	P
<i>Octolasion tyrtaeum</i>	P	P	P
Endemic genus/species			
<i>Curgiona</i>	A	A	P
<i>Kotegeharia</i>	A	A	P
<i>Mallehulla</i>	A	A	P
<i>Priodochaeta</i>	A	A	P
<i>Karmiella</i>	A	A	P
<i>Troyia</i>	A	A	P
<i>Comarodrilus</i>	A	A	P
<i>Chaetocotoides</i>	A	A	P
<i>Parryodrilus</i>	A	A	P
<i>Dashiella</i>	A	A	P
<i>Moniligaster</i>	A	A	P
<i>Celeriella</i>	A	A	P
<i>Lampito</i>	A	A	P
<i>Travoscolides</i>	A	A	P
<i>Wahoscolex</i>	A	A	P
<i>Tonoscolex</i>	A	P	A
<i>Kanchuria</i>	A	P	A
<i>Perionyx</i> (4 species)	P	A	A
<i>Eutyphoeus</i> (2 species)	P	A	A
<i>Plutellus</i> (1 species)	P	A	A

Table 2: Density and biomass of earthworms (m⁻²) in different areas in the Indian mountainous region (blank fields indicate non-availability of data)

Land use/ ecosystems	Diversity	Density (number of individuals m ⁻²)	Average/ range of biomass (wet weight, g)	Trends	References
Pine woodland, Meghalaya		9-32	4,4-9,4		Reddy and Alfred, Shilling, (1978)
Mixed woodland, H. P.		30-118			Julka and Mukherjee-Solan, (1984)
Humid Tropical deciduous forest, A. P. (Eastern Ghats)	5 Species: <i>Amyntas alexandri</i> , <i>A. diffringens</i> , <i>Metaphire posthuma</i> , <i>M. houletti</i> , <i>Dichogaster</i> sp.	315(28-281 in different micro sites within forest ecosystem)		Two peaks, one during & the other at the end of the rainy season	Reddy (1987)
Shifting agriculture at Nangpoh in North-Eastern India	3 species: <i>Megascolides antrophytes</i> , <i>Drawida assamensis</i> , <i>Nelosclex strigosus</i>	68		Peak density immediately after rainy season	Mishra and Ramakrishnan, 1988
Shifting agriculture at Shilling in north-eastern India	5 Species: <i>Amyntas diffringens</i> , <i>Drawida assamensis</i> , <i>Eutyphoeus festvus</i> , <i>Nelosclex phase strigosus</i> , <i>Tonoscolex horatii</i>	447 in cropping phase, 50 in fallow		Peak density during rainy season except <i>Amyntas</i> in winter	Bhadauria and Ramakrishnan, 1989
Pasture Kumaon Himalaya	3 species: <i>Amyntas diffringens</i> , <i>Amyntas alexandri</i> , <i>Eisenia fetida</i>	139	25	Peak density at the end of rainy season	Kaushal and Bisht, 1994
Cultivated soil Kumaon Himalaya	One species: <i>Amyntas alexandri</i>	58		Peak density during rainy season	Kaushal et al. (1995)
A village landscape in Garhwal Himalaya	8 species: <i>Amyntas alexandri</i> , <i>Bimastos parvus</i> , <i>Metaphire anomala</i> , <i>Metaphire birmanica</i> , <i>Drawida nepalensis</i> , <i>Perionyx excavatus</i> , <i>Lennogaster pusillus</i> , <i>Octochaetona beatrix</i>	Traditional agro forestry system: 147; Traditional pure crop system: 132; Moderately degraded natural forest: 63; Rehabilitated Agricultural land: 63; Abandoned agricultural land: 27; Rehabilitated forest land: 8; Highly degraded forest land: 5		All species except <i>B. parvus</i> showed a strong effect of season on population size, with the highest abundance and biomass values observed during rainy season. Only <i>D. nepalensis</i> and <i>M. birmanica</i> were a little bit abundant during dry season in traditional agro forestry system	Bhadauria et al. (2012)
Village landscape in Kumaon Himalaya	8 Species: <i>Bimastos parvus</i> , <i>Octolasion tytaeum</i> , <i>Octochaetona beatrix</i> , <i>Amyntas corticis</i> , <i>Eutyphoeus festvus</i> , <i>E. naniianus</i> , <i>E. waltonii</i> , <i>Drawida</i> sp.	Climax forest: 526, Mixed forest: 309, 5-year-old pine: 287, 40-year-old pine: 940, grassland :353		Peak density during rainy season	Bhadauria and Ramakrishnan, 2000
Village landscape in Garhwal Himalaya	7 Species: <i>Amyntas corticis</i> , <i>Drawida nepalensis</i> , <i>Allobophora parva</i> , <i>Eutyphoeus pharpiangianus</i> , <i>Octochaetona beatrix</i> , <i>Perionyx</i> sp., <i>Lennogaster pusillus</i>	108-247 in forests, 89-235 in agro-ecosystems		Abundance in pine forest- higher but diversity lower compared to oak forest	Sinha et al. (2003)
Nandadevi Biosphere Reserve in Garhwal Himalaya	8 species: <i>Lennogaster pusillus</i> , <i>Metaphire houletti</i> , <i>M. anomala</i> , <i>Ocneroдрilus occidentalis</i> , <i>Dendrodrilus rubidus</i> , <i>Aporrectodea calliginosa</i> , <i>Amyntas corticis</i> , <i>Drawida nepalensis</i>	Lower elevation: Home garden- 100, pine forest-5-10, oak forest: 5, irrigated agriculture-5, Rainfed agriculture: 5-25 Higher elevation-Home garden:10-150 Alpine pasture:5-25; Alpine forest: 10-15 10-262(mean ~ 109)Pontoscolex corethrurus-dominant 72% of total density, <i>Kanchuria</i> spp-1-12%, <i>Metaphire houletti</i> -3.6%	9- 100	Peak density at the end of rainy season	Maikhuri et al., 2008
Rubber plantations in Tripura in the north-eastern Himalaya	20 species: <i>Glossoscolecidae</i> - <i>Pontoscolex corethrurus</i> , <i>Octochaetidae</i> - <i>Eutyphoeus gigas</i> , <i>E.gammiei</i> , <i>E. comillahnus</i> , <i>E. assamaensis</i> , <i>E. festvus</i> , <i>Eutyphoeus</i> spp., <i>Dichogaster bolau</i> , <i>D.affinis</i> , <i>Lennogaster chittagongensis</i> , <i>Octochaetona beatrix</i> , <i>Megascolecidae</i> - <i>Metaphire houletti</i> , <i>Perionyx</i> spp., <i>Kanchuria sumerianus</i> , <i>Kanchuria</i> sp 1 & sp.2, <i>Moniligastridae</i> - <i>Drawida nepalensis</i> , <i>Drawida</i> sp-1, <i>Drawida</i> sp-2, <i>Gordiodrilus elegans</i>		9-100 (mean 43.4) <i>Pontoscolex corethrurus</i> - 61.5%Of total biomass		Chaudhuri et al. (2008)

Table 2: Density and biomass of earthworms (m⁻²) in different areas in the Indian mountainous region (blank fields indicate non-availability of data)

Land use/ ecosystems	Diversity	Density (number of individuals m ⁻²)	Average/ range of biomass (wet weight, g)	Trends	References
Nilgiri, Biosphere Reserve, Kerala area in the Western Ghats	Fourteen species: <i>Dichogaster affinis</i> , <i>Drawida modesta</i> , <i>D. banwelli impertusa</i> , <i>D. ghatensis</i> , <i>D. grandis</i> , <i>Glyphidrilus amandalei</i> , <i>Haplochetalla sp.</i> , <i>Lampito mauritii</i> , <i>Megascolex insignis</i> , <i>M. triangularis</i> , <i>Octochaetona beatrix</i> , <i>Paryodrilus lavellei</i> , <i>Plutellus variabilis</i> , <i>Pontoscolex corethrus</i>	Natural forests: 159; Forest plantations: 350; Palms: 258; Croplands: 640; Fallows: 124			Chandrashekara et al. (2008)
Nilgiri Biosphere reserve; Karnataka area in the Western Ghats	13 species: <i>Drawida somavarapatna</i> , <i>Drawida modesta</i> , <i>Drawida lakir</i> , <i>Drawida pellicuda</i> , <i>Lampito mauritii</i> , <i>Metaphire holoutii</i> , <i>Megascolex curgenis</i> , <i>Megascolex felisceta</i> , <i>Amyntas corticis</i> , <i>Octochaetoides castellanes</i> , <i>Pontoscolex corethrus</i> , <i>Dendrodrilus rubidus</i> , <i>Gargia narayani</i>	Natural forests: 477; Forest plantations: 303; Cardamom plantations: 443; Croplands: 595; Fallows: 343			
A segment of Western Ghats		14-825		Earthworms mainly exhibited temporal variability while some other soil organisms were affected more by management practices	Rossi and Blanchart (2005)

Table 3: Ecological category, feeding habit, and habitat and size relationship of eight species of earthworms in Tripura (Chaudhuri et al., 2008; Chaudhuri and Bhattacharjee, 2009)

Species	Family	Size(mm) L, length; B, breadth	Color	Feeding habit	Distributi on pattern	Ecological category	Soiltemp (°C) (pH: 4.4-5.2)	Soilmoisture (g %v)	Soil pH	Soil organic matter(g %)
<i>Pontoscolex corethrus</i>	Glossoscolecidae	L = 72-100B = 4-5	Lightly pigmented	Geophagous	Exotic	Topsoil	19-32	10-29	4.5-4.8	1.7-2.4
<i>Drawida papillifer</i>	Moniligastridae	L = 45-90B = 3-4	Deeply pigmented	Phytophagous	Endemic	Endogaeic	22-32	10-29	4.5-4.7	1.6-2.4
<i>Drawida daassamensis</i>	Moniligastridae	L = 60-80B = 4-5	Lightly pigmented	Phytophagous	Endemic	Topsoil endogaeic	21-31	10-29	4.4-5.2	1.5-2.0
<i>Metaphire houlleti</i>	Megascolecidae	L = 100-160B = 3-6	Deeply pigmented dorsally	Phytophagous	Endemic	Epianecic	21-28	14-20	4.4-5.2	1.6-2.0
<i>Eutyphoeus comillahnus</i>	Octochaetidae	L = 70-135B = 2-4	Lightly pigmented	Geophagous	Endemic	Mesohumic endogaeic	21-27	17-18	4.5-4.6	1.7-2.0
<i>Dichogaster affinis</i>	Octochaetidae	L = 35-42B = 1-2	Moderately pigmented	Phytophagous	Exotic	Epigeic	21-28	17-21	4.8-5.2	1.8-2.0
<i>Octochaetona beatrix</i>	Octochaetidae	L = 60-120B = 4-5	Lightly pigmented	Geophagous	Endemic	Subsoil endogaeic	24-28	11..5-14	4.7-4.8	1.6-2.0
<i>Lenngaster chittagongensis</i>	Octochaetidae	L = 40-50B = 1.5-2.5	Lightly pigmented	Phytophagous	Endemic	Topsoil endogaeic	24-30	16-18	4.7-4.8	1.5-2.0

Ghats (Dash and Senapati, 1985; Dash and Dash, 2008; Dash et al., 2009). *Haplochetalla* spp is widespread in laterite and red soils of the Western Ghats.

Studies in different land-uses in the state of Karnataka in south India showed species composition of earthworm communities in natural forests closer to that in plantations (coffee, Acacia and cardamom plantations) and paddy fields but radically different from that in grasslands. *Pontoscolex corethrurus*, an endogeic exotic species, was found in all land-uses (Kale et al., 2008, Chandrashekara et al., 2008). Studies carried out in Eastern Ghat region also showed variation in species composition between land uses but a species like *Lampito mauritii* persisted in all land uses (Dash and Patra, 1977; Senapati and Dash, 1981; Mishra and Dash, 1984; Senapati et al., 2005).

Population and biomass

Data available on earthworm population diversity, density, biomass in different mountainous regions of India are summarized in Table 2. Earthworm density and biomass are influenced by a whole range of abiotic and biotic factors. A high level of environmental heterogeneity and variation in land use/management practices in mountains may result in huge variation in biodiversity within an Agroclimatic region. *Metaphire anomala*, *Metaphire houlletii*, *Ocneroдрilus occidentalis*, *Dendrodrilus rubidus* and *Aporrectordea calliginosa* occurred in the Nanda Devi Biosphere Reserve, a relatively cool area with lower degree of anthropogenic pressures due to legal protection but not in Hariyali landscape, a comparatively warmer area faced to more intense human disturbances in the absence of any legal protection. *Allobophora parva*, *Eutyophoeus pharapingianus*, *Octochaetona beatrix* and *Perionyx* spp. occurred in the latter




Figure 1: Agroclimatic regions of India (from Julka and Paliwal, 2005)




Figure 2: Land use change and earthworm community structure in north-eastern Himalaya (Bhadauria and Ramakrishnan, 2005)

area but not in the former area (Sinha et al., 2003; Maikhuri et al., 2008). In areas with high intensity of agricultural land use, earthworm community comprises *Amynthus diffringens*, *Amynthus alexandri* and *Eisenia fetida* (Kaushal and Bisht, 1994; Kaushal et al., 1995), species rarely found in traditional landscapes with less intensively land uses. The latter do differ in terms of structure and composition of earthworm communities. *Bimastos parvus* and *Octolasion tyrtaeum* were sampled from a village landscape in Almora district (Bhadauria and Ramakrishnan, 2000) but not from a similar landscape in Chamoli district (Sinha et al., 2003). In shifting agricultural landscapes in the Eastern Himalaya, *Drawida assamensis* and *Nelloscolex strigosus* occurred across an elevation gradient while *Megascolides astrophytes* was confined to lower elevations and *Amynthus diffringens* and *Tonoscolex horai* to higher elevations (Mishra and Ramakrishnan, 1988; Bhadauria and Ramakrishnan, 1989). Total number of species in cultural landscapes did not vary much (6-8) but certain ecosystem types/patches may be quite poor (e.g., wet paddy fields, Cedrus forests subject to intensive disturbances) or quite rich (e.g., home gardens, rainfed agriculture) in terms of number of species (Maikhuri et al., 2008; Bhadauria et al., 2012).

Earthworms could be a sensitive indicator of environmental quality, e.g., presence of *Ramiellona wilsoni* is indicative of pristine tropical montane forests in Mexico (Negrete-Yankelevich et al., 2007) but necessarily not in all situations. Earthworm population was not significantly influenced by intensity of management in grasslands in Ireland (Curry et al., 2008) and tropical rain forest ecozone in the Western Ghats of India (Rossi and Blanchart, 2005). Density and biomass are likely to be more sensitive to environmental changes and land management practices than species richness and diversity. In

Table 4: Earthworm species richness in Easternghat land uses (Based on Mishra and Dash, 1984; Senapati et al., 2005)

S.no.	Species	Land Uses				
		Natural Forest	Disturbed	Shifting Cultivation 8-year old fallow	Cropping phase	Eucalyptus plantation
1.	<i>Drawida calebi</i>	+	+	-	-	-
2.	<i>Drawida willsi</i>	+	-	-	-	-
3.	<i>Eutyphoeus incommodes</i>	+	-	+	+	-
4.	<i>E.waltoni</i>	-	-	-	+	-
5.	<i>Eutyphoeus</i> spp.	+	-	-	+	-
6.	<i>Lampito mauritii</i>	+	+	-	-	+
7.	<i>Lenogaster dashi</i>	-	-	-	-	+
8.	<i>Lenogaster pusillus</i>	+	-	+	+	-
9.	<i>Ocnerodrilus occidentalis</i>	+	+	-	-	-
10.	<i>Octochaetona surensis</i>	+	+	-	-	-
11.	<i>Pellogaster bengalensis</i>	+	+	+	+	-
12.	<i>Ramiella bishambari</i>	+	+	+	-	-

the Nanda Devi Biosphere Reserve, earthworm abundance declined with decline in temperature (*i.e.*, increase in elevation) and was sensitive to both season and land use management, with significant interactions between species, season and management practices. Flooded paddy systems had the lowest and the home gardens the highest species diversity as well as abundance (Maikhuri et al., 2005). Earthworm abundance in the village landscapes at lower elevations was higher (89-940 individuals per m²) compared to that at higher elevations (5-150 per m²) (Table 2). In western/central Himalayan region and Tripura in north-eastern Himalaya, endogeic and endogeic-anecic dominate. In rubber plantations in Tripura established after 1962, 15 species are endogeic and only five species are epi-anecic. Further, rubber plantations gave way to exotics like *Pontoscolex corethrurus* (Chaudhuri et al., 2008; Chaudhuri and Bhattacharjee, 2009).

In Nilgiri Biosphere Reserve, 14 species of earthworms were recorded, with 2-8 species occurring in different land use types and total earthworm abundance in the range of 124-560 per m². Occurrence of endogeic *Parryodrilus lavelee* and *Pontoscolex corethrurus* in almost all land uses including degraded lands suggests that these species may have a potential for rapid restoration of soil fertility in degraded lands (Chandrashekara et al., 2008). In most studies, population size has been estimated in terms of numerical abundance and not in terms of biomass.

Table 3 gives ecological category, feeding habit, and habitat and size relationship of eight species, out of 20 species of earthworms of rubber plantations raised in undulating areas in Tripura (Chaudhuri et al., 2008, Chaudhuri and Bhattacharjee, 2009). This type of study will be useful for Himalayan and other regions to identify the suitable species for land use management. Functional attributes and ecological strategies of different earthworm species need to be worked out to optimize the contributions of earthworms to ecosystem services and resilience of agroecosystems.

Functional attributes: Bioturbation activity

Bioturbation refers to the biological reworking of soil and sediments, and its importance was first highlighted by Charles Darwin (1881). Bioturbation is now recognized as an archetypal example of 'ecosystem engineering', modifying geochemical gradients, redistributing food resources and

microbes in soil column. Bioturbation played a key role in the evolution of metazoan life at the end of the Precambrian Era (Muys et al., 2003). Earthworm casts contain more water soluble aggregates and higher nutrient concentrations than the surrounding soils. Soils with earthworms drain 4 to 10 times faster than soils without earthworms (Guild, 1952, 1955; Low, 1955; Dash and Patra, 1979; Petersen and Luxton, 1982; Bhadauria and Ramakrishnan, 1989; Bhadauria et al., 1997).

Bhadauria and Ramakrishnan (1991) estimated cast production at a rate of 20 tons, 35 tons, 40 tons per hectare per year in a 5-year-old pine forest, a 35-year-old pine forest, and a sacred grove (close to climax vegetation), respectively, in the north-eastern hill region of India. Chaudhuri et al. (2008) estimated cast production at a rate of 2.51 ton per ha per year in rubber plantations in Tripura in the north-eastern India. These rates of cast production are substantially lower than the rates of 77-141 tons hectare per year reported in temperate/tropical ecosystems (Satchell, 1967; Dash and Patra 1979). Estimation of cast production in different land uses would indicate the functional role of earthworms but such studies covering different land-uses and eco-regions in India are lacking.

Land use change and earthworms

Bhadauria and Ramakrishnan (2005) worked on earthworm community structure in relation to land use change in shifting agricultural landscapes in the north-eastern Himalaya (Fig. 2), Chaudhuri et al. (2008, 2009) in rubber plantations in Tripura, Chandrashekara et al. (2008) in Nilgiri Biosphere, Senapati et al. (1994 and 2002) in tea gardens in south India and Mishra and Dash (1984) (Table 4), Dash and Senapati (1991); Behera et al. (1999) and Senapati et al. (2005) in Orissa in south-east India.

Based on these studies, some generalizations can be made. Conversion of natural forests to shifting agriculture and plantations results in some loss of earthworm species richness together with changes in composition of soil fauna community structure and function. Exotic and native species coexist in natural and derived (managed) ecosystems but exotics are less frequent in primary forests. Endopolyhumics in primary forests and Endomesohumics in derived ecosystems (man-managed) dominate the earthworm community structure in north-east India. However in central Himalayas,

endomesohumic earthworms dominated irrespective of land use changes from primary forest to grassland and fallows. Land use change from forest to agro ecosystem favoured endomesohumic possibly due to high input of farmyard manure in settled agroecosystems in central/western Himalaya. Senapati *et al.* (2005) observed proliferation of termite populations following land use intensification in tea plantations and Senapati *et al.* (1994) and Senapati (1997) suggested that termite-earthworm biomass ratio can be used as a sensitive index to land use change. This hypothesis is yet to be tested in wide variety of ecosystems.

CONCLUSIONS

Conversion of natural forest results in decline of earthworm species richness and abundance. Organic inputs in the form of manure and crop residues in the derived systems help in restoring earthworm fauna. Termites may proliferate at the expense of earthworms in these derived systems. Periodically inoculation of endogeic and endo-anecic earthworms is likely to restore soil quality in degraded ecosystems. The selection of earthworm species should centre on the naturalized species in a particular site irrespective of whether the species is native or exotic.

ACKNOWLEDGEMENT

This publication arises from our association with and support from Tropical Soil Biology and Fertility Institute of CIAT (TSBF-CIAT), TSBF-UNEP-GEF programme on belowground biodiversity and MMSEA-SLM network of United Nations University Institute of Sustainability and Peace (UNU-ISP), Tokyo.

REFERENCES

- Behera, B. Giri, S., Dash, N. C., Sahu, J. and Senapati, B. K. 1999. Earthworm bioindication of forest land use pattern in western Orissa. *Indian Forester*. **125**: 272-281.
- Bhadoria, T. and Ramakrishnan, P. S. 1991. Population dynamics of earthworms and their activities in forest ecosystems of north-east India. *J. Tropical Ecology*. **7**: 305-318.
- Bhadoria, T. and Ramakrishnan, P. S. 1989. Earthworm population dynamics and contribution to nutrient cycling during cropping and fallow phases of shifting agriculture (Jhum) in northeast India. *J. Applied Ecology*. **26**: 505-520.
- Bhadoria, T., Ramakrishnan, P. S. and Srivastava, K. N. 2000. Diversity and distribution of endemic and exotic earthworms in natural and regenerating ecosystems in the central Himalaya, India. *Soil Biology and Biochemistry*. **32**: 2045-2054.
- Bhadoria, T. and Ramakrishnan, P. S. 2005. Impact of land use-land cover change on earthworm community structure in India. pp. 57-65. In: Ramakrishnan, P. S., Saxena, K. G., Swift, M. I., Rao, K. S. and Maikhuri, R. K. (Editors) *Soil Biodiversity, Ecological processes and Landscape Management*. Oxford and IBH Publishers, New Delhi, India.
- Bhadoria, T., Kumar, P., Kumar, R., Maikhuri, R. K., Rao, K. S. and Saxena, K. G. 2012. Earthworm populations in a traditional village landscape in Central Himalaya, India. *Applied Soil Ecology*. **53**: 83-93.
- Bhadoria, T., Ramakrishnan, P. S. and Srivastava, K. N. 1997. Population dynamics of earthworms during crop rotation under rain fed agriculture in central Himalayas, India. *Applied Soil Ecology*. **6**: 205-215.
- Chandrashekhara, U. M., Balasundaram, M., Sankaran, Sujatha, M. P., Varma, R. V., Senapati, B. K. and Sehgal, M. 2008. Conservation and Sustainable Management of Below-ground Biodiversity in the Kerala part of Nilgiri Biosphere Reserve-phase I, Final Technical Report, Kerala Forest Research Institute, Thissur, India. p.119.
- Chaudhuri, P. S. and Bhattacharjee, S. 2009. Reproductive biology (cocoon structure, incubation period, emergence pattern and fecundity) of eight tropical earthworm species of rubber plantations in laboratory cultures. *J. Tropical Ecology* (In press)
- Chaudhuri, P. S., Nath, S. and Paliwal, R. 2008. Earthworm population of rubber plantations (*Hevea brasiliensis*) in Tripura India. *Tropical Ecology*. **49**: 225-234
- Choudhuri, P. S. 2009. Personal communication.
- Darwin, C. 1881. The formation of vegetable mould through the action of worms with observations on their habits. John Murray, London
- Dash, M. C. 1978. The role of earthworms in the decomposer system, Pages 399-409, In: Singh, J. S. and Gopal, B. (Eds) *Glimpses of Ecology*, International Scientific Publishers, Jaipur, India
- Dash, M. C. 2012. Charles Darwin's Plough. Tools for Vermitechnology. I. K. International Publishing House Pvt. Ltd. New Delhi. p. 185.
- Dash, M. C. and Patra, U. C. 1977. Density, biomass and energy budget of a tropical earthworm population from Orissa, India. *Review of Ecology and Biology of Soil*. **14**: 461-471.
- Dash, M. C. and Patra, U. C. 1979. Worm cast production and nitrogen contribution to soil by a tropical earthworm population from a grassland site in Orissa, India. *Review of Ecology and Biology of Soil*. **16**: 79-87.
- Dash, M. C. and Senapati, B. K. 1991. Impact of Eucalyptus litter on earthworm activity. In: *Advances in Management and Conservation of Soil Fauna*. Veeresh, G.K., Rajgopal, D. and Viraktamaath, C.A. (Editors), Oxford and IBH Publication Company Private Limited, New Delhi. pp. 349-355.
- Dash, M. C. and Senapati, B. K. 1985. Vermitechnology potentiality of Indian earthworms for vermicomposting and vermifeed. In: Mishra, H.K. and Kapur, K. (Editors) *Proceedings of National Seminar on Current Trends in Soil Biology*. Haryana Agriculture University, Hissar, India. pp 61-69.
- Dash, M. C., Saxena, K. G. and Giri, S. 2009. Vermitechnology for Wasteland Reclamation, Plant Productivity and Composting: A Review in Indian Context. *International J. Ecology and Environmental Sciences*. **35(2-3)**: 163-185.
- Dash, M. C. and Dash, S. P. 2008. Conservation and sustainable management of belowground biodiversity: A review on the functional role of soil fauna in Indian ecosystems with particular reference to earthworms. *International J. Ecology and Environmental Science*. **34(3)**: 223-243.
- Guild, W. J. Mc.L. 1952. Variations in earthworm numbers within field population. *J. Animal Ecology*. **21**: 169
- Guild, W. J. Mc.L. 1955. Earthworms and Soil Structure. pp. 83-98, In: Kevan, D.K. Mc.E. (Editor) *Soil Zoology*, Butterworth's, London.
- Julka, J. M. 2008. Personal communication.
- Julka, J. M. and Mukherjee, R. N. 1984. Mass migration of earthworm in the Simla hill. Page 347-351, In: *Proceedings of workshop on High Altitude Entomology and Wildlife*, Zoological Survey of India, Calcutta.
- Julka, J. M. and Paliwal, R. 2005a. Diversity and Biogeography of Indian Earthworms, In: *proceedings of National Level Workshop on Vermitechnology Transfer to NSS programme officers*, Department of Biochemistry and Zoology, Kongunadu Arts and Science College, Coimbatore, India. pp. 5-21.

- Julka, J. M. and Paliwal, R. 2005b.** Distribution of earthworms in different agro-climatic regions of India. In: Soil Biodiversity, Ecological Processes and Landscape Management. Edited by P.S. Ramakrishnan, K.G. Saxena, M.J. Swift, K.S. Rao and R.K. Maikhuri. Oxford and IBH Publishing Company Private Limited, New Delhi. pp. 3-13
- Kale, R. D. and Dinesh, M. S. 2005.** Earthworm: Components of Soil Biota, In: Proceedings of National Workshop, Kerala Forest Research Institute, Peechi, Kerala, India. pp. 313-330.
- Kareiva, P. and Marvier, M. 2003.** Conserving biodiversity coldspots. *American Scientist*. **91**: 344-351.
- Kaushal, B. R. and Bisht, S. P. S. 1994.** Population dynamics of the earthworm *Amyntas alexandri* (Annelida: Megascolecidae) in Kumaon Himalayan pasture soil. *Bio. and Fertility of Soils*. **17**: 9-13.
- Kaushal, B. R., Bisht, S. P. S. and Kalia, S. 1995.** Population dynamics of earthworm *Amyntas alexandri* (Megascolecidae: Annelida) in cultivated soils of the Kumaon Himalayas. *Applied Soil Biology*. **2**: 125-130.
- Low, A. J. 1955.** Improvement in the structure of soils under leys. *J. Soil Science*. **6**: 179-199.
- Maikhuri, R. K., Rawat, L. S., Negi, G. S., Dangwal, D., Chandra, A. and Rao, K. S. 2005.** Macro fauna in Nanda Devi Biosphere Reserve. In: The Proceedings of the National Workshop on Conservation and Sustainable Management of Belowground Biodiversity, Kerala Forest Research Institute, Peechi, India, pp. 422-431 (unpublished document).
- Maikhuri, R. K., Dangwal, D., Misra, S., Rawat L. S., Srivastava, A., Singh, K., Mustafa, Negi, C. S., Chamoli, K. P. and Rao, K. S. 2008.** Conservation and sustainable management of belowground diversity in the Nanda Devi Biosphere Reserve, final technical report, phase I, G. B. Pant Institute of Himalayan Environment and Development, Garhwal unit, Uttarakhand, India. p. 150.
- Mishra, K. C. and Ramakrishnan, P. S. 1988.** Earthworm population dynamics in different Jhum fallows developed after slash and burn agriculture in north-eastern India. *Proceedings Indian Academy of Sciences (Animal Sciences)*. **97**: 309-318.
- Mishra, P. C. and Dash, M. C. 1984.** Population dynamics and respiratory metabolism of earthworm population in subtropical woodland of western Orissa, India. *Tropical Ecology*. **25**:103-116.
- Muys, B., Beckers, G., Nachtergale, L., Lust, N., Merck, R. and Granval, P. 2003.** Medium term evaluation of a forest soil restoration trial combining tree species change, fertilization, and earthworm introduction. *Pedobiologia*. **47**: 772-783.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B. and Kent, J. 2000.** Biodiversity hotspots for conservation priorities. *Nature*. **403**: 853-858.
- Negrete Yankelevich, S., Fragoso, C., Newton, A. C. and Heal, O. W. 2007.** Successional changes in soil, litter and macroinvertebrate parameters following selective logging in a Mexican Cloud Forest. *Applied Soil Ecology*. **35**: 340-355.
- Nijhawan, S. D. and Kanwar, J. S. 1952.** Physicochemical properties of earthworm castings and their effect on productivity of the soil. *Indian Journal of Agricultural Sciences*. **22**: 357-373
- Petersen and Luxton, M. 1982.** A Comparative analysis of soil fauna populations and their role in decomposition process. *Oikos*. **39**: 287-388.
- Ramakrishnan, P. S. Saxena, K. G., Swift, M. J., Rao, K. S. and Maikhuri, R. K. 2005.** Soil Biodiversity, Ecological Processes and Landscape Management. Oxford and IBH Publishing Company Private Limited, New Delhi. p.302.
- Reddy, M. V. 1987.** Seasonal structure of earthworm (*Pheretima* sp.) Populations in humid tropical deciduous woodland. In: Bonvicini Pagliai, M and Omodeo, P (Editors) On Earthworms, Selected Symposia and Monographs, U. Z. I., Mucchi, Modena. pp. 191-197
- Reddy, M. V. and Alfred, J. R. B. 1978.** Some observations on the earthworm population and their biomass in a sub-tropical pine forest soil. In: Edwards, C. A. and Veeresh, G. K. (Editors) Soil Biology and Ecology in India. University of Agricultural Sciences, Bangalore. pp. 78-82
- Rossi, J. P. and Blanchart, E. 2005.** Seasonal and land-use induced variations of soil macro fauna composition in the Western Ghats, southern India. *Soil Biology and Biochemistry*. **37**: 1093-1104.
- Satchell, J. E. 1967.** Lumbricidae. In: Burges, A and Raw, F.B. (Editors), Soil Biology, Academic Press, London. pp. 259-322
- Senapati, B. K. and Dash, M. C. 1981.** Effect of grazing on the elements of production in vegetation and Oligochaeta of a tropical pasture. *Review of Ecology and Biology of Soil*. **18**: 487-505.
- Senapati, B. K., Julka, J. M. and Lavelle, P. 2005.** Impacts of Land use-land cover change on soil fauna in south and south-east India. In: Ramakrishnan, P. S., Saxena, K. G., Swift, M. J., Rao, K. S. and Maikhuri, R. K. (Editors), Soil Biodiversity, Ecological Processes and Landscape Management, Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi. pp. 67-75
- Senapati, B. K., Lavelle, P., Panigrahi, P. K., Giri, S. and Brown, G. G. 2002.** Restoring soil fertility and enhancing productivity in Indian tea plantations with earthworm and organic fertilizers. International Technical Workshop on Biological Management of Soil Ecosystems for Sustainable Agriculture. *Brazilian Agriculture Research Corporation, Brazil*. **182**: 172-190.
- Senapati, B. K., Panigrahi, P. K. and Lavelle, P. 1994.** Macrofaunal status and restoration strategy in degraded soil under intensive tea cultivation in India. In: Transactions of 15th International Soil Science Congress. *Acapulco and Mexico*. **4(a)**: 64-75
- Senapati, B. K. 1997.** Innovations in Soil management through earthworm technology: A case study in tea agroecosystems, In: Modern Approaches and Innovations in Soil Management. Bagyaraj, D. J., Verma, A., Khanna, K. K. and Kheri, H.K. (Editors), Rastogi Publications, Meerut, pp. 282-300.
- Sinha, B., Bhadauria, T., Ramakrishnan, P. S., Saxena, K. G. and Maikhuri, R. K. 2003.** Impact of landscape modification on earthworm diversity and abundance in the Hariyali sacred landscape, Garhwal Himalaya. *Pedobiologia*. **47**: 357-370.
- State of Forest Report, Government of India, 2005 and 2009.